



cancer.org | 1.800.227.2345

Urostomy Guide

Urostomy surgery is needed when the bladder isn't working the way it should. There are 4 major bladder problems that may be treated with a urostomy:

- [Bladder cancer](#)
- Damage to the nerves that control the bladder (called neurogenic bladder disease)
- Birth defects
- Chronic inflammation of the bladder

For the thousands of people who have serious bladder diseases, a urostomy can be the start of a new and healthier life. If you have had a chronic (long-term) problem or a life-threatening disease like bladder cancer, you can look forward to feeling better after you recover from urostomy surgery. You can also look forward to returning to most, if not all of the activities you enjoyed in the past.

This guide will help you better understand urostomy – what it is, why it's needed, how it affects the normal urinary system, and the changes it can bring to a person's life.

- [What Is a Urostomy?](#)
- [Types of Urostomies and Pouching Systems](#)
- [Caring for a Urostomy](#)

What Is a Urostomy?

A urostomy is an opening in the belly (abdominal wall) that's made during surgery. It re-

directs urine away from a bladder that's diseased, has been injured, or isn't working as it should. The bladder is either bypassed or removed. (Surgery to remove the bladder is called a *cystectomy*.) After this surgery, urine is passed out of the body through an opening on the belly called a *stoma*. A Wound Ostomy Continence nurse (WOCN or WOC nurse) or the surgeon will figure out the best location for your stoma. (A WOC nurse is a specially trained registered nurse who takes care of and teaches ostomy patients. This nurse may also be called an *ostomy nurse*.)

- [Why would you need a urostomy?](#)
- [Getting ready for surgery](#)

The stoma will look pink to red and will be moist and shiny. The shape will be round to oval, and it will shrink over time after surgery. Some stomas may stick out a little, while others are flat against the skin.

You won't be able to start or stop urine coming out through the stoma, so you will likely need a pouch to collect the urine as it comes out. The stoma has no nerve endings, so it's not a source of pain or discomfort. It is just a change in the way urine comes out of your body.

A urostomy is not a disease, but it's usually needed because of a disease or other problem. A urostomy causes a change in [the way your body works](#)¹. It surgically changes the way urine comes out of your body.

Why would you need a urostomy?

Urostomy surgery is done when certain diseases and conditions cause serious bladder problems.

If there's cancer in the bladder, part of the treatment may be to remove all or part of the bladder and divert or detour the urine by doing a urostomy.

Some people find it easier to manage a urostomy than a defective bladder. Bladder problems often mean that people can't control the flow of urine – they are incontinent. This can be stressful, and the constant wetting may cause skin problems.

Some children are born with a defect in the urinary tract that causes urine to back up into the kidneys. This leads to chronic or repeat infections. For these children, a urostomy may be life-saving. Surgery can make a pathway through which the urine may travel easily, without any blockages (or obstructions) that cause it to back up. This allows the kidneys to work their best.

Getting ready for surgery

As you get ready for abdominal surgery, your doctor and/or ostomy nurse will explain the process and look at your abdomen to find the best location for the stoma. You may be asked to wear a sample pouch to make sure that the place chosen is on the flattest possible surface and that you are comfortable in all positions. If you have any hobbies, activities, or habits that might be affected by where the pouch is placed, talk to the doctor or the ostomy nurse.

Talking with someone who has gone through the same experience can help, too. Look for an ostomy visitor program in your area - an ostomy nurse can help with this. This is a program that matches you with a volunteer from the United Ostomy Association of America who has a urostomy and has adapted well to the changes that it brings. They can answer many of your questions about day-to-day life. You may also find it helpful to take part in an ostomy support group. A support group allows you to share your feelings and ask questions as you recover and adjust to your urostomy. You can also share your story with others who may be helped by your experience.

Hyperlinks

1. www.cancer.org/cancer/types/bladder-cancer/about/what-is-bladder-cancer.html

References

In its original form this document was written by the United Ostomy Association of America (1962-2019). It has since been modified and updated by the American Cancer Society using the following sources.

Berti-Hearn L, Elliott B. Urostomy care: A guide for home care clinicians. *Home Healthcare Now*. 2019; 37(5):248-255.

Hollister. *Urostomy: What's right for me*. Accessed at https://www.hollister.com/~//media/files/pdfs-for-download/ostomy-educational-booklets/hol_os_urostomy-whats-right_na_907603.pdf on October 2, 2019.

United Ostomy Association of America (UOAA). *New ostomy patient guide: Colostomy, ileostomy, urostomy, continent diversion*. Accessed at https://www.ostomy.org/wp-content/uploads/2018/05/All-In-One-New-Patient-Guide_2018.pdf on October 2, 2019.

United Ostomy Association of America (UOAA). *Urostomy guide*. Accessed at <https://www.ostomy.org/wp-content/uploads/2018/03/UrostomyGuide.pdf> on October 2, 2019.

Last Revised: October 16, 2019

Types of Urostomies and Pouching Systems

There are 2 basic options for urostomy surgery, often called **urinary diversion**. The 2 types are named for how the pathway that allows urine to pass from the body is diverted (re-routed or changed) during the procedure.

- **Incontinent diversion**, sometimes called a **standard or conventional** urostomy
 - **Continent diversion**, also called a **continent urostomy** or **continent urinary reservoir**
-
- [Incontinent diversion \(standard or conventional urostomy\)](#)
 - [Continent diversion \(continent urostomy or urinary reservoir\)](#)
 - [Types of pouching systems](#)
 - [Choosing a pouching system for standard urostomy](#)
-

Neurologic
dysfunction

Birth defects


stop) drainage

adaptable to night drainage

Chronic
inflammation
of bladder

For this type of urostomy, an **ileal conduit** is made. An ileal conduit is a small pouch that holds urine. It's surgically created from a small piece of bowel (intestine).

To make an ileal conduit, a 6- to 8- inch piece of the lower part of the small intestine (called the **ileum**) is cut out near where it attaches to the large intestine (colon). The rest of the ileum is then re-connected to the large intestine. The ureters are detached from the bladder and attached to the piece of ileum that was removed. The bladder may or may not be removed. The far end of the piece of ileum is brought through the front of the abdomen wall to form the stoma. The other end is sewn closed to make a pocket that holds the urine inside the body. This surgery is a life-long change. The urine output cannot be controlled and a collection pouch will have to be worn at all times.



The small intestine produces mucus naturally. The segment of intestine that was used to form the ileal conduit or urine pocket will keep making mucus, too. The mucus will collect in the pouch along with the urine. This is normal.

Continent diversion (continent urostomy or urinary reservoir)

For this type of urostomy, the surgeon also makes an internal pouch. In this case, the pouch has valves to keep the urine from backing up the ureters into the kidneys. There's also a valve to keep the urine in the pouch until it's removed. The pouch must be drained 4 or 5 times a day with a thin, flexible tube called a *catheter*. The doctor or ostomy nurse will teach you how to drain the pouch and help you make a schedule to do it.

Types of continent diversion urostomies


There are several different ways of creating a continent diversion urostomy, each using different parts to create the storage pouch, valves, and outlet.

Kock pouch: the pouch, valves, and outlet are made from the end of the small intestine (terminal ileum).

Indiana pouch: the pouch is made from the large intestine (ascending colon). The outlet is made from the end of the small intestine (terminal ileum), and the natural ileocecal valve is used.

Mitrofanoff procedure: this type of diversion can have several different variations. The pouch can be made from the bladder, large or small intestines, or a combination. The outlet is made from the appendix, a fallopian tube, or part of the ureter.

Ileal neobladder: the pouch is made from the small intestine (ileum). For a neobladder, there is no ostomy, and urine is passed through the urethra. You won't have the urge to urinate, so a schedule is needed.



Types of pouching systems

Pouches come in many styles and sizes, and an ostomy nurse can help you choose the best one for your situation and lifestyle. They all have a collection pouch to collect the urinary drainage and an adhesive barrier that sticks to your skin (called a flange, skin barrier, or wafer) and a collection pouch. There are 2 main types of systems available.

- One-piece pouches have both a pouch and skin barrier attached together in the same unit. When the pouch is removed, the barrier also comes off.
- Two-piece systems have a pouch and a separate skin barrier. When the pouch is taken off, the barrier stays in place.



Two-Piece Pouch

While you are in the hospital after your surgery, an ostomy nurse will talk to you about your personal needs and help you pick a pouching system. For some, this pouch is the type worn for a lifetime. Others will need to change to a new or different type of pouching system later on if they gain weight, grow (in the case of a child), or for other reasons. Do not keep using a recommended pouch if it's not working the way you need it to. Try different types until you find one that feels good, works well, and is easy to use.



Choosing a pouching system for standard urostomy

You will need a pouching system to collect urine that comes out of the stoma. It must have a barrier that protects the skin around it.

It's best to talk with an ostomy nurse or someone who has experience in this area when you're trying out your first pouching system. There should be someone in the hospital who can get you started with equipment and instructions after surgery.

As you're getting ready to leave the hospital, be sure you get referrals to an ostomy nurse, an ostomy clinic, an ostomy manufacturer, or a chapter of the United Ostomy Associations of America. Even if you must go out of town to get such help, it's worthwhile, as you want to get a good start and avoid mistakes. Even with help, you may have to try different types or brands of pouching systems to find the one that best suits you.

The size of the stoma, abdominal firmness and shape, the location of the stoma, scars and folds near the stoma, and your height and weight all must be considered. Special changes may have to be made for stomas near the hipbone, waistline, groin, or scars. Some companies have custom-made products to fit unusual situations.

A good pouching system should:

- Stay secure with a good leak-proof seal that lasts for 3 to 7 days
- Be odor-resistant
- Protect the skin around the stoma
- Be nearly invisible under clothing
- Be easy to put on and take off
- Allow you to shower or bathe with the pouch on, if you wish to do so

Factors that can affect the pouching system seal

The pouching system must stick to your skin. It's important to change it before it loosens or leaks. The length of time the pouch stays sealed to the skin depends on many things, such as proper fitting, the weather, skin condition, scars, weight changes, diet, activity, and the shape of your body near the stoma. Here are some other things that may affect how long a pouch sticks:

- Sweating will shorten the number of days you can wear a pouch. Body heat, added to outside temperature, will cause skin barriers to loosen more quickly than usual.
- Moist, oily skin may reduce wearing time.
- Weight changes will affect how long you can wear a pouch. Weight gained or lost after urostomy surgery can change the shape of your abdomen. You may need an entirely different system.
- Physical activities may affect wearing time. Swimming, very strenuous sports, or anything that makes you sweat may shorten wearing time.

Using a night drainage system

At night the bottom of the pouch can be connected to a night drainage system which carries urine away from the stoma while you sleep. This lets you sleep undisturbed. Gravity drains the urine to a bedside container. This keeps your pouch from getting so full and heavy that it pulls away from your body. It also protects your stoma and skin against the build-up of strong urine.

When connecting the pouch to the bedside drainage bag, leave a small amount of urine in the pouch before attaching the tubing. This keeps you from setting up a vacuum in the system. The bedside container should be vented and the tubing should be secured at the top of the bag or bottle with no more than one inch of it going down into the bag or bottle. If the urine backs up into the tubing it will stop draining. The drainage container can be hung on the side of the bed, or put in another container on the floor. A small wastebasket may be used for this purpose. Clean the tubing and bedside drainage collector every morning. Pour about 2 ounces of a vinegar-and-water solution into the emptied pouch that's attached to the night drainage system. (Use 1 part of white vinegar to 3 parts of water.) Let it run out of the pouch, through the tubing, and into the collection container. Then pour it out and hang up the night drainage tubing so it can air dry.

Night Drainage System

References

In its original form this document was written by the United Ostomy Association of America (1962-2019). It has since been modified and updated by the American Cancer Society using the following sources.

Berti-Hearn L, Elliott B. Urostomy care: A guide for home care clinicians. *Home Healthcare Now*. 2019; 37(5):248-255.

Hollister. *Urostomy: What's right for me*. Accessed at https://www.hollister.com/~media/files/pdfs-for-download/ostomy-educational-booklets/hol_os_urostomy-whats-right_na_907603.pdf on October 2, 2019.

United Ostomy Association of America (UOAA). *New ostomy patient guide: Colostomy, ileostomy, urostomy, continent diversion*. Accessed at https://www.ostomy.org/wp-content/uploads/2018/05/All-In-One-New-Patient-Guide_2018.pdf on October 2, 2019.

United Ostomy Association of America (UOAA). *Urostomy guide*. Accessed at <https://www.ostomy.org/wp-content/uploads/2018/03/UrostomyGuide.pdf> on October 2, 2019.

Last Revised: October 16, 2019

Caring for a Urostomy

As you learn more, you may have questions about caring for your urostomy. This section has many tips that come from others. But remember, no two people are alike. There's no one right answer, only suggestions and ideas for you to try.

- [Emptying and changing the pouching system](#)
- [Protecting the skin around the stoma](#)
- [Shaving hair under the pouch](#)
- [Spots of blood on the stoma](#)
- [Eating restrictions](#)
- [Bathing](#)
- [When you should call the doctor](#)
- [Ordering and storing ostomy supplies](#)

Emptying and changing the pouching system

You'll be taught how to change and empty your pouching system before you leave the hospital. You don't have to use sterile supplies. For instance, facial tissue, toilet paper, or paper towels can be used to clean around the stoma instead of sterile gauze pads.

Emptying the pouch

The urostomy pouch has a drain valve at the bottom, so it can be emptied as needed. Since bacteria grow quickly in urine, it's important to empty the pouch often, at regular times. A lot of urine in the bag also could damage the pouch seal. It's a good idea to empty your pouch when it's about 1/3 to 1/2 full.

During the day most people need to empty the pouch about as often as they used the bathroom before they had urostomy surgery or other bladder problems – for many people, this might mean every 2 to 4 hours, or more often if you drink a lot of fluids. Children may have to empty more often because their pouches are smaller. Follow these steps:

- Sit as far back on the toilet as you can.
- Place a small strip of toilet paper in the toilet to decrease splashing.
Hold the bottom of the pouch up to reduce splashing.

- Close the valve.
- Dry the end of the valve with toilet paper.

At night a piece of flexible tubing can be attached to the drain valve on your pouch. This allows the urine to flow into a bigger bedside bag or drainage bottle while you sleep. Many people find a bedside drainage unit better than getting up during the night to empty the pouch. (This is discussed further in "Using a night drainage system.")

Changing the pouch

Different pouching systems are made to last different lengths of time. Some are changed every day, some every 3 days or so, and some just once a week. It depends on type of pouch you use.

Your pouch should be changed on a schedule that fits your routine. And it's best to have a regular changing schedule so problems don't develop. In other words, don't wait for it to leak to change it. It may be easier to change the pouch in the morning before you eat or drink anything. If this isn't a good time for you, try to wait at least 1 to 2 hours after drinking fluids so that urine doesn't get on your skin and supplies while you change the pouch.

You'll want to find the best position for putting on your pouch – sitting, standing, or lying down. The position you use should allow you to see your stoma and be easy for you to be in when making a change. Some people stand facing the toilet so the urine dripping from the stoma falls in the toilet. When changing while sitting in a wheelchair, it's helpful to slide your buttocks toward the front of the chair and lean back a little. Using a mirror may help you center the pouch over the stoma.

Before changing your pouch, clean your hands well and put all your supplies on a clean surface. Clean pouches decrease the chances of germs (bacteria) getting into your urinary system. Bacteria can multiply quickly even in the tiniest drop of urine. These germs may travel up the ureters and cause a kidney infection. Bacteria can also cause foul-smelling urine. Some people find it helpful to use rolled tissues, paper towels, or tampons to absorb dribbling urine from the stoma.

You don't have to use sterile supplies – the stoma and nearby skin are clean, but not sterile. But keeping all your supplies clean and organized will save you time and money. Always have at least 2 complete pouches, one on your body and one ready for your next change.

At first you may find that changing your pouch can take up to 30 minutes. With time and

from the skin.

Clean the skin around the stoma with water. This can be done in the shower or tub. Pat the skin dry before putting on the skin barrier or pouch.

Watch for sensitivities and allergies to the adhesive, skin barrier, paste, tape, or pouch material. They can develop after weeks, months, or even years of using a product because you can become sensitized over time. If your skin is irritated only where the plastic pouch touches it, you might try a pouch cover or a different brand of pouch. A stoma nurse can offer ideas if needed. Pouch covers are available from supply manufacturers, or you can make your own. You may have to **test different products to see how your skin reacts to them.**

Shaving hair under the pouch

Having a lot of hair around the stoma can make it hard to get the skin barrier to stick well and may cause pain when you remove it. Shaving with a razor or trimming hair with

- When the food you eat is used by the body, it leaves a mineral residue called ash. This ash can be either acidic or basic (alkaline) depending on the type of food. Most fruits and vegetables give an alkalinized ash and tend to make the urine alkaline. Meats and cereals usually lead to an acidic ash residue.
- Unless you are told otherwise, your urine should be kept in an acid state. Drink cranberry juice in place of orange juice or other citrus juices (these tend to make the urine more alkaline), and take vitamin C daily (check with your doctor first). Some of the acid ash foods include: most meats, breads and cereals, cheese, corn, cranberries, eggs, macaroni, nuts, pasta, prunes, fish, and poultry.
- Urinary crystals on the stoma or skin are caused by alkaline urine. The crystals look like white, gritty particles. They may lead to stoma irritation and/or bleeding. Proper cleaning, keeping your urine acidic, and careful fitting of the skin barrier will help prevent urinary crystals.
- To help reduce urinary crystals, you can make a vinegar compress and apply it to the stoma for a few minutes when the pouch is changed. To do this, soak a bath cloth or small towel in a mixture of equal parts of water and white vinegar and hold the moist cloth on the stoma.

- you notice several times when emptying it)
- Bleeding where the stoma meets the skin
 - Bad skin irritation or deep sores (ulcers)
 - An unusual change in your stoma size or color
 - Fever or strong urine odor (may be a sign of a kidney infection)

Ordering and storing ostomy supplies

Supplies may be ordered from a mail order company, from a medical supply store, or from a local pharmacy. If you want to order supplies online. For additional help ordering, try contacting the product manufacturer, check your local phone book's business pages, or check the Internet (try searching for "ostomy supplies").

- To order pouches, skin barriers, and other ostomy products, you'll need both the **manufacturer's name and the product numbers**
- Talk with your ostomy nurse about reputable dealers who can supply you with what you need
- Before ordering your supplies, check with your health insurance to be sure that they work with the company in question
- Compare prices when using mail order and the Internet (remember to include shipping costs)
- Keep all your supplies together on a shelf, in a drawer, or in a box in a dry area away from moisture and hot or cold temperatures.
- Order supplies a few weeks before you expect them to run out to allow enough time for delivery. **DO NOT** stockpile supplies – they can be ruined by moisture and temperature changes.

References

In its original form this document was written by the United Ostomy Association of America (1962-2019). It has since been modified and updated by the American Cancer Society using the following sources.

Berti-Hearn L, Elliott B. Urostomy care: A guide for home care clinicians. *Home Healthcare Now*. 2019; 37(5):248-255.

Hollister. *Urostomy: What's right for me*. Accessed at https://www.hollister.com/~media/files/pdfs-for-download/ostomy-educational-booklets/hol_os_urostomy-whats-right_na_907603.pdf on October 2, 2019.

United Ostomy Association of America (UOAA). *New ostomy patient guide: Colostomy, ileostomy, urostomy, continent diversion*. Accessed at https://www.ostomy.org/wp-content/uploads/2018/05/All-In-One-New-Patient-Guide_2018.pdf on October 2, 2019.

United Ostomy Association of America (UOAA). *Urostomy guide*. Accessed at <https://www.ostomy.org/wp-content/uploads/2018/03/UrostomyGuide.pdf> on October 2, 2019.

Last Revised: October 16, 2019

Written by

The American Cancer Society medical and editorial content team
(<https://www.cancer.org/cancer/acs-medical-content-and-news-staff.html>)

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as editors and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy (www.cancer.org/about-us/policies/content-usage.html).

cancer.org | 1.800.227.2345