



[cancer.org](https://www.cancer.org) | 1.800.227.2345

Waldenstrom Macroglobulinemia Early Detection, Diagnosis, and Staging

Know the signs and symptoms of Waldenstrom macroglobulinemia. Find out how Waldenstrom macroglobulinemia is tested for, diagnosed, and staged.

Detection and Diagnosis

Catching cancer early often allows for more treatment options. Some early cancers may have signs and symptoms that can be noticed, but that is not always the case.

- [Can Waldenstrom Macroglobulinemia Be Found Early?](#)
- [Signs and Symptoms of Waldenstrom Macroglobulinemia](#)
- [How Is Waldenstrom Macroglobulinemia Diagnosed?](#)

Stages and Outlook (Prognosis)

After a cancer diagnosis, staging provides important information about the extent of cancer in the body and anticipated response to treatment.

- [How Is Waldenstrom Macroglobulinemia Staged?](#)
- [Survival R rgi for moldenstrom Macroglobulinemia](#)

Can Waldenstrom Macroglobulinemia Be Found Early?

Waldenstrom macroglobulinemia (WM) is not common, and at this time there are no widely recommended screening tests to look for this disease in people without symptoms.

Still, many cases of WM are found early, either when people go to the doctor because of symptoms they are having, or when they have blood tests done for other reasons. The best way to find this cancer early is to see your doctor if you have signs or symptoms that might be caused by this disease.

Hyperlinks

1. www.cancer.org/cancer/types/waldenstrom-macroglobulinemia/references.html

References

Last Revised: July 19, 2018

Castillo JJ. Plasma cell disorders. *Prim Care*. 2016; 43:667-691. doi: 10.1016/j.pop.2016.07.002. Epub 2016 Oct 14.

Rajkumar SV, Dispenzieri A. Chapter 104: Multiple myeloma and related disorders. In: Niederhuber JE, Armitage JO, Dorshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 5th ed. Philadelphia, Pa. Elsevier: 2014.

[See all references for Waldenstrom Macroglobulinemia](#)

Last Revised: July 19, 2018

Signs and Symptoms of Waldenstrom Macroglobulinemia

- [Common symptoms of WM](#)
- [Less common signs and symptoms of WM](#)

Sometimes, Waldenstrom macroglobulinemia (WM) isn't causing any symptoms when it's first found. Instead, it's found when the person has blood tests done for some other

from the buildup of the abnormal protein.

Loss of appetite: Some people with WM lose their appetite.

Fever, sweats, weight loss: WM, like other lymphomas, can cause fevers (without an infection), drenching night sweats, and weight loss (without trying). These are called **B symptoms**.

Neuropathy: In some people with WM, the abnormal antibody can attack and damage nerves outside the brain. This can lead to numbness or a painful “pins and needles” sensation in the feet and legs, which is called neuropathy.

Less common signs and symptoms of WM

Enlarged lymph nodes: These usually appear as lumps under the skin around the neck, in the groin, or in the armpits. Enlarged lymph nodes are usually about 1 or 2 inches (2.5-5 cm) across. They are seen less often in WM than in most other lymphomas.

Swollen abdomen (belly): WM can sometimes make the spleen or liver bigger, making the belly look swollen. In the upper part of the abdomen, the liver is on the right and the spleen on the left. When the spleen gets larger, it can press on the stomach, which makes people feel full when they eat even a small amount.

Circulation system symptoms: In hyperviscosity syndrome, the thickened blood causes poor brain circulation, leading to problems like headache, confusion, and dizziness. It can also cause symptoms like those seen with a stroke, including slurred speech and weakness on one side of the body. Patients with these symptoms should contact their doctor right away.

Abnormal bleeding: High levels of abnormal antibody can damage blood vessels, which can lead to problems like nosebleeds and bleeding gums.

Vision problems:

Heart problems: High levels of the M protein can damage heart tissue directly or through the development of amyloidosis, in which the protein builds up in the heart muscle. This weakens the heart, affecting its ability to pump blood. In addition, because the blood of people with WM is thicker than normal, their hearts have to work harder to pump blood throughout the body. This strain can wear down the heart muscle, leading to a condition called **congestive heart failure**. Symptoms can include heart palpitations, feeling tired and weak, cough, shortness of breath, rapid weight gain, and swelling in the feet and legs.

Infections: The high levels of abnormal antibody in WM can slow the body's normal antibody production. This makes it harder for the body to fight infections.

Digestive symptoms: In some people with WM, the buildup of the M protein in the intestines can lead to problems such as diarrhea, poor absorption of vitamins, or gastrointestinal bleeding (seen as blood in the stools or dark stools).

Sensitivity to cold: In people with cryoglobulins, exposure to cold temperatures can lead to pain, itching, a bluish color, or even sores on the tip of the nose, ears, fingers, or toes due to reduced blood flow to these areas.

Hyperlinks

1. www.cancer.org/cancer/types/non-hodgkin-lymphoma.html
2. www.cancer.org/cancer/types/waldenstrom-macroglobulinemia/references.html

References

Castillo JJ. Plasma cell disorders. *Prim Care*. 2016; 43:667-691. doi: 10.1016/j.pop.2016.07.002. Epub 2016 Oct 14.

Rajkumar SV, Dispenzieri A. Chapter 104: Multiple myeloma and related disorders. In: Niederhuber JE, Armitage JO, Dorshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 5th ed. Philadelphia, Pa. Elsevier: 2014.

[See all references for Waldenstrom Macroglobulinemia](#)

Last Revised: July 19, 2018

How Is Waldenstrom Macroglobulinemia Diagnosed?

- [Medical history and physical exam](#)
- [Lab tests](#)
- [Biopsies](#)
- [Lab tests on biopsy specimens](#)
- [Imaging tests](#)

Waldenstrom macroglobulinemia (WM) is often found when a person goes to see their doctor because of symptoms they are having, or because they just don't feel well and go in for a checkup. Sometimes it's found in people without symptoms when they have blood tests done for some other reason.

If signs or symptoms suggest that a person might have WM, exams and tests will be done to be sure. The most important tests will look for abnormal proteins in the blood and abnormal cells in the bone marrow. Because WM is a type of [lymphoma](#)¹, like other lymphomas it can invade the bone marrow, lymph nodes, and other organs.

Medical history and physical exam

Your doctor will want to get a thorough medical history, including information about your

symptoms, possible [risk factors](#)², family history, and other medical conditions.

Next, the doctor will examine you, paying special attention to your lymph nodes and other areas of your body that might be involved, including the eyes, nerves, spleen, and liver. The doctor might also look for signs of infection, which can cause many of the same symptoms.

If the doctor suspects that WM (or another type of lymphoma) might be causing your symptoms, the next step probably will be to order blood tests. You might also be referred to a hematologist, a doctor who specializes in diseases of the blood, or an oncologist, a doctor who specializes in cancer.

Lab tests

WM might be suspected if your doctor finds you have low blood cell counts or unusual protein levels on blood tests. If so, your doctor may order a blood test called **serum protein electrophoresis** to find out what the abnormal proteins are. It is usually only after these tests are done that a biopsy of either the bone marrow or a lymph node is considered.

Blood cell counts

The complete blood count (CBC) is a test that measures the levels of red blood cells, white blood cells, and platelets. If lymphoma cells occupy too much of the bone marrow, these blood levels may be low.

Immunoglobulin levels

This test measures the levels of the different antibodies (immunoglobulins) in the blood – IgA, IgE, IgG, and IgM – to see if any are abnormally high or low. obBl 9 gs (ta0 0od)Tj 0 g 1 0 0 1 72

Finding a monoclonal IgM antibody in the blood is needed to diagnose WM. This abnormal protein in WM is known by many different names, including **monoclonal immunoglobulin M, IgM protein, IgM spike, IgM paraprotein, M protein, and M-spike**. High levels of other types of monoclonal immunoglobulins, like IgA or IgG, are seen in different disorders (like [multiple myeloma](#)³ and some other lymphomas).

Sometimes pieces of the IgM protein are excreted by the kidneys into the urine. These proteins can be detected with a test called **urine protein electrophoresis** (or UPEP).

Viscosity

Viscosity is a measure of how thick the blood is. If the IgM level is too high, the blood will become thick (viscous) and can't flow freely (think about pouring honey compared to pouring water).

Cryocrit

This test measures the blood levels of cryoglobulins (proteins that clump together in cool temperatures and can block blood vessels).

Cold agglutinins

Cold agglutinins are antibodies that attack and kill red blood cells, especially at cooler temperatures. These dead cells can then build up and block blood vessels. A blood test can be used to detect these antibodies.

Beta-2 microglobulin (2M)

This test measures another protein made by the cancer cells in WM. This protein itself doesn't cause any problems, but it's a useful indicator of a patient's prognosis (outlook). High levels of 2M are linked with a worse outlook.

Biopsies

The symptoms of WM and non-Hodgkin lymphoma (NHL) are not distinctive enough for a doctor to know for certain if a person has one of them, based on symptoms alone. Most symptoms can also be caused by non-cancerous problems like infections or by other kinds of cancers. Blood tests can help point to the correct diagnosis, but a biopsy (removing samples of affected tissue to look at under a microscope) is the only way to be sure. Several types of biopsies might be used.

Bone marrow aspiration and biopsy

This is the most important type of biopsy for WM, and is needed to confirm the diagnosis. It can be done at the doctor's office or at the hospital.

The bone marrow aspiration and biopsy are usually done at the same time. The samples are taken from the back of the pelvic (hip) bone, although in some cases they may be taken from the sternum (breast bone) or other bones.

- In bone marrow **aspiration**, you lie on a table (either on your side or on your belly). The doctor cleans the skin over the hip and then numbs the area and the surface of the bone by injecting a local anesthetic. This may briefly sting or burn. A thin, hollow needle is then inserted into the bone, and a syringe is used to suck out a small amount of liquid bone marrow. Even with the anesthetic, most patients still have some brief pain when the marrow is removed.
- A bone marrow **biopsy** is usually done just after the aspiration. A small piece of bone and marrow is removed with a slightly larger needle that is pushed down into the bone. This may also cause some brief pain.

Once the biopsy is done, pressure is applied to the site to help stop any bleeding. There will be some soreness in the biopsy area when the numbing medicine wears off. Most patients can go home right after the procedure.

The bone marrow samples are then sent to a lab, where they are tested to see if they have lymphoma cells (see below). For a diagnosis of WM, at least 10% of the cells in the bone marrow must be lymphoplasmacytoid lymphoma cells.

Fine needle aspiration (FNA) biopsy

In an FNA biopsy, the doctor uses a very thin, hollow needle with a syringe to withdraw a small amount of tissue from a tumor or lymph node. This type of biopsy is useful for sampling lymph nodes to see if they are enlarged because of cancer or an infection. FNA can help diagnose some lymphomas, but WM is usually diagnosed with a bone marrow biopsy.

For an FNA on an enlarged node near the surface of the body, the doctor can aim the needle while feeling the node. If the enlarged node (or tumor) is deep inside the body, the needle can be guided while it is seen on a computed tomography (CT) scan or

Flow cytometry

In this test, cells are treated with special man-made antibodies. Each antibody sticks only to certain types of cells. The cells are then passed in front of a laser beam. If the cells now have antibodies attached to them, the laser will make them give off light, which is measured and analyzed by a computer.

This is the most common test for **immunophenotyping** – classifying lymphoma cells according to the proteins (antigens) on their surfaces. Different types of lymphocytes have different antigens on their surface. These antigens also change as each cell matures.

This test can help show if a lymph node is swollen because of lymphoma, some other cancer, or a non-cancerous disease. It has become very important in helping doctors determine the exact type of lymphoma so they can select the best treatment.

Cytogenetics

Doctors use this technique to look at the chromosomes (long strands of DNA) inside lymphoma cells. Cells (usually from the bone marrow) are first grown in the lab. Then the chromosomes are stained and looked at closely. Because it takes time for the cells to start dividing, this test can take a few weeks.

In some lymphomas, the cells may have too many chromosomes, too few chromosomes, missing parts of chromosomes (called deletions), or other abnormalities. These changes can help identify the type of lymphoma.

Molecular genetic tests

Molecular tests such as fluorescent in situ hybridization (FISH) and polymerase chain reaction (PCR) are not usually needed to diagnose WM, but they are sometimes used to diagnose other types of NHL. These tests look at the cells' DNA without having to grow the cells in the lab first. The tests can give results in less time than cytogenetics and can be done on cells from different sources (like lymph nodes, blood, and bone marrow). They are generally used to look for specific chromosome or gene changes, not just any change.

More testing information

See [Testing Biopsy and Cytology Specimens for Cancer⁴](#) to learn more about tests used to diagnose cancer and what the results can tell you.

A [PET scan](#)⁷ can be helpful in spotting small collections of cancer cells. It is even more valuable when combined with a CT scan (PET/CT scan).

PET scans also can help tell if an enlarged lymph node contains lymphoma or not. It can help spot small areas that might be lymphoma, even if the area looks normal on a CT scan. These tests can be used to tell if a lymphoma is responding to treatment. They can also be used after treatment to help decide whether an enlarged lymph node still contains lymphoma or is merely scar tissue.

Hyperlinks

1. www.cancer.org/cancer/types/lymphoma.html
 2. www.cancer.org/cancer/types/waldenstrom-macroglobulinemia/causes-risks-prevention/risk-factors.html
 3. www.cancer.org/cancer/types/multiple-myeloma.html
 4. www.cancer.org/cancer/diagnosis-staging/tests/biopsy-and-cytology-tests.html
 5. www.cancer.org/cancer/diagnosis-staging/tests/imaging-tests/ct-scan-for-cancer.html
-

beta-2 microglobulin (2M). People with lower levels of IgM and 2M tend to do better than those with higher levels. People with WM who are older, are anemic (based on a low blood hemoglobin level), or have a low blood platelet count tend to have a poorer outlook.

Experts have used these factors to develop a system that helps predict prognosis (outlook) for patients with WM. It is called the **International Prognostic Scoring System for Waldenstrom Macroglobulinemia** (ISSWM). This system takes into account the factors that seem to predict a poorer outcome, such as:

- Older than 65
- Blood hemoglobin level 11.5 g/dL or less
- Platelet count 100,000/mcL or less
- Beta-2 microglobulin more than 3 mg/L
- Monoclonal IgM level more than 7 g/dL

Except for age, each of these factors is worth a single point. The points are added to make a score, which is used to divide patients into 3 risk groups:

- The low-risk group includes patients 65 or younger who have no more than 1 point.
- The intermediate-risk group includes those who are older than 65 with 2 or fewer points, and those younger than 65 who have 2 points.
- The high-risk group includes those of any age who have at least 3 points.

These groups can be used to help predict survival (discussed in more detail in [Survival Rates for Waldenstrom Macroglobulinemia](#)).

Hyperlinks

Morel P, Duhamel A, Gobbi P, et al. International prognostic scoring system for Waldenstrom macroglobulinemia. *Blood*. 2009;113:4163–4170.

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology: Waldenstrom's macroglobulinemia/Lymphoplasmacytic lymphoma. V.1.2018. Accessed at www.nccn.org/professionals/physician_gls/pdf/waldenstroms.pdf on June 21, 2018.

[See all references for Waldenstrom Macroglobulinemia](#)

Last Revised: July 19, 2018

Survival Rates for Waldenstrom Macroglobulinemia

- [What is a 5-year survival rate?](#)
- [Cancer survival rates don't tell the whole story](#)

Survival rates are often used by doctors as a way of discussing a person's outlook.

Survival rates tell you what percentage of people with the same [type](#)¹ and stage of cancer are still alive a certain length of time (usually 5 years) after they were diagnosed. These numbers can't tell you how long you will live, but they may help give you a better understanding about how likely it is that your treatment will be successful.

What is a 5-year survival rate?

Statistics on the outlook for a certain type and stage of cancer are often given as 5-year survival rates, but many people live longer – often much longer – than 5 years. The 5-year survival rate is the percentage of people who live at least 5 years after being diagnosed with cancer. For example, a 5-year survival rate of 12% means that 12 out of 100 people diagnosed with cancer will live at least 5 years after diagnosis.

diagnosed.

Relative survival rates

ISSWM risk group 5-year survival rate

Low	87%
Intermediate	68%
High	36%

Median survival

Median survival is another way to look at survival. It is the length of time at which half of the patients in a group are still alive, and half have died. By definition, half of the patients live longer than the median survival. The group that developed the ISSWM used data from WM patients diagnosed and treated before January 2002 and found the following:

ISSWM risk group Median survival*

Low	12 years
Intermediate	8 years
High	3.5 years

*Median survival is measured from the point that treatment is started.

In the last decade (2001-2010), the median overall survival for all WM groups has improved to just over 8 years compared to 6 years in the previous decade (1991-2000).

Hyperlinks

1. www.cancer.org/cancer/types/waldenstrom-macroglobulinemia/about/what-is-wm.html
2. doi.org/10.1111/bjh.13264
3. www.cancer.org/cancer/types/waldenstrom-macroglobulinemia/references.html

References

Castillo JJ, Olszewski AJ, Cronin AM, Hunter ZR, Treon SP. Survival trends in

Waldenström macroglobulinemia: An analysis of the Surveillance, Epidemiology and End Results database. *Blood*. 2014;123:3999–4000.

Castillo, J. J., Olszewski, A. J., Kanan, S. , Meid, K. , Hunter, Z. R. and Treon, S. P. (2015), Overall survival and competing risks of death in patients with Waldenström macroglobulinaemia: an analysis of the Surveillance, Epidemiology and End Results database. *Br J Haematol*, 169: 81-89. doi:[10.1111/bjh.13264](https://doi.org/10.1111/bjh.13264)*Blood*. 2014;123:3999–400035Tj 0 g ETO

Questions to Ask About Waldenstrom Macroglobulinemia

- Do you recommend starting treatment now or waiting until later on?

When deciding on a treatment plan

- What are my [treatment options](#)¹?
- What do you recommend and why?
- Is a [stem cell transplant](#)² an option for me? What are the pros and cons of this treatment?
- How much experience do you have treating this type of cancer?
- Should I get a [second opinion](#)³? How do I do that? Can you recommend someone?
- What would the goal of the treatment be (to reduce symptoms, lower IgM levels, etc.)?
- How quickly do we need to decide on treatment?
- What should I do to be ready for treatment?
- How long will treatment last? What will it be like? Where will it be done?
- What risks or side effects are there to the treatments you suggest? Are there things I can do to reduce these side effects?
- How might treatment affect my daily activities? Can I still work full time?
- What are the chances the cancer will recur (come back) with these treatment plans?
- What will we do if the treatment doesn't work or if the cancer recurs?
- What if I have transportation problems getting to and from treatment?

During treatment

Once treatment begins, you'll need to know what to expect and what to look for. Not all of these questions may apply to you, but asking the ones that do may be helpful.

- How will we know if the treatment is working?
- Is there anything I can do to help manage side effects?
- What symptoms or side effects should I tell you about right away?
- How can I reach you on nights, holidays, or weekends?
- Do I need to change what I eat during treatment?
- Are there any limits on what I can do?
- Can I exercise during treatment? If so, what kind should I do, and how often?
- Can you suggest a mental health professional I can see if I start to feel

overwhelmed, depressed, or distressed?

- What if I need social support during treatment because my family lives far away?

After treatment

- Do I need a special diet after treatment?
- Are there any limits on what I can do?
- What other symptoms should I watch for?
- What kind of exercise should I do now?
- What type of follow-up will I need after treatment?
- How often will I need to have follow-up exams and imaging tests?
- Will I need any blood tests?
- How will we know if the cancer has come back? What should I watch for?
- What will my options be if the cancer comes back?

Along with these sample questions, be sure to write down some of your own. For instance, you might want more information about recovery times. Or you may want to ask about available [clinical trials](#)⁴.

Keep in mind that doctors aren't the only ones who can give you information. Other health care professionals, such as nurses and social workers, can answer some of your questions. To find out more about speaking with your health care team, see [The Doctor-Patient Relationship](#)⁵.

Hyperlinks

1. www.cancer.org/cancer/types/waldenstrom-macroglobulinemia/treating.html
2. www.cancer.org/cancer/types/waldenstrom-macroglobulinemia/treating/stem-cell-transplantation.html
3. www.cancer.org/cancer/managing-cancer/finding-care/seeking-a-second-opinion.html
4. www.cancer.org/cancer/managing-cancer/making-treatment-decisions/clinical-trials.html
5. www.cancer.org/cancer/managing-cancer/finding-care/the-doctor-patient-relationship.html

